




Gymnosperms vs. Angiosperms




Gametophyte/Sporophyte Cycle



Angiosperm Reproduction



Seed Development




Monocot vs. Dicot

Copyright © The McGraw-Hill Companies, Inc. Permission is granted to reproduce or display.

	Seed	Root	Stem	Leaf	Flower
Monocots	One cotyledon in seed	Root system and phloem in a ring	Vascular bundles scattered in stem	Leaf veins form a parallel pattern	Flower parts in threes or multiples of three
Eudicots	Two cotyledons in seed	Root phloem between arms of xylem	Vascular bundles in a distinct ring	Leaf veins form a net pattern	Flower parts in fours or fives and their multiples

Types of Fruits



Plant Transport

Short-distance

occurs by osmosis (movement of free water through aquaporins)

Long-distance

bulk flow using:
translocation (H^+ gradient) through phloem (sugar)
transpirational pull (cohesion) through xylem (fluid/minerals)

Plant Hormones

Auxins

distribution of auxins → phototropism
enhances apical dominance (vertical)
promotes fruit growth

Cytokinins

stimulates cytokinesis & cell division
delays senescence (aging)

Gibberellins

cause stem elongation, fruit growth, & seed germination
auxin + gibberellins = fruit development

Brassinosteroids

induce cell elongation & division in stem segments, slow leaf abscission

Abscisic Acid (ABA)

antagonizes growth hormones, inhibits germination, used to withstand drought
must be washed/exposed to light/cold to deactivate

Ethylene Gas

produced in response to stress during fruit ripening, apoptosis, & exposure to auxin