

### Enzymes

**Enzyme** Biological molecules (**proteins**) that act as **catalysts** and **help complex reactions**.

are specific to their substrates and each enzyme has its own optimum pH

**Substrate** Material upon which an enzyme acts.

**Enzyme inhibition** important in normal metabolism for control of pathways.

### Reversible inhibitor

**competitive** (Raises  $K_m$  only) Same size and shape with the substrate inhibition

**noncompetitive** (Lowers  $V_{max}$  only) inhibitor doesn't mind whether there is a substrate or not. but when the inhibitor binds, it switches off catalysis.

**uncompetitive** (Lowers  $V_{max}$  and  $K_m$ ) the inhibitor can ONLY be on the surface of the enzyme if the substrate is there.


**Irreversible inhibitor** acts by reacting with the enzyme protein, usually at the active site(substrate site), to permanently block activity.

$V_{max}$  is the maximum rate of an enzyme catalysed reaction i.e. when the enzyme is saturated by the substrate.


$K_m$  is measure of how easily the enzyme can be saturated by the substrate.

$K_m$  and  $V_{max}$  are constant for a given temperature and pH and are used to characterise enzymes. They can be used to identify types of inhibitors i.e. competitive, non-competitive and uncompetitive.


### Allosteric site




### Normal binding



### Enzyme kinetics



### Reversible inhibition




### Enzyme in human digestion

Three main types of enzymes in human digestion:

- Amylases** break down carbohydrates
  - Example: salivary amylase
  - Substrate: starch Product: maltose
  - Source: mouth (salivary glands)
  - Optimum pH: 7-7.8
- Proteases** break down polypeptides
  - Example: pepsin
  - Substrate: polypeptides Product: amino acids
  - Source: stomach
  - Optimum pH: 2
- Lipases** break down fats and lipids
  - Example: pancreatic lipase
  - Substrate: triglycerides Product: fatty acids & glycerol
  - Source: pancreas, delivered into small intestine
  - Optimum pH: 7.2 - 7.5

### with co-enzyme



### CoFactor

In the case of **metal ion cofactors**

**Stabilise** the structure, not directly involved in the chemistry

**Ca++** with some proteinases

**part of substrate** **Mg++-ATP** with some kinases

**part of active site** **Zn++** in alcohol dehydrogenase

### Organic cofactors

**coenzymes** they do come on and off like other substrates (NAD+)

**prosthetic groups** a cofactor that forms a permanent part of the enzyme's active site.

Doesn't come on and off in a catalytic cycle (FAD, PLP)


### NAD+ is both a coenzyme and a substrate

Why use coenzyme in one case and not in the other case?

Metabolic point: NAD is one of a range of cofactor substances that is present in small concentration

They turn over and over again to process a large amount of substances

Pathway is to process large amount with tiny amount of coenzyme



By **rhettbro**  
[cheatography.com/rhettbro/](https://cheatography.com/rhettbro/)

Published 12th April, 2021.  
Last updated 14th April, 2021.  
Page 1 of 2.

Sponsored by **Readable.com**  
Measure your website readability!  
<https://readable.com>